arXiv:math/0102184v1 [math.GT] 23 Feb 2001

SOME HOMOLOGICAL INVARIANTS OF MAPPING CLASS GROUP OF A 3-DIMENSIONAL HANDLEBODY

SUSUMU HIROSE

ABSTRACT. We show that, if $g \geq 2$, the virtual cohomological dimension of the mapping class group of a 3-dimensional handlebody of genus g is equal to $4g-5$ and the Euler number of it is equal to 0.

1. INTRODUCTION

A genus g *handlebody* H_g is an oriented 3-manifold which is constructed from 3-ball with attaching g 1-handles. The *mapping class group* $\mathcal{H}_g$ of H_g is defined as the group of isotopy classes of orientation-preserving diffeomorphisms of H_g. This group $\mathcal{H}_g$ is a subgroup of the mapping class group $\mathcal{M}_g$ of a surface ∂H_g, that is, $\mathcal{M}_g = \pi_0(\mathrm{Diff}^+(\partial \mathrm{H_g}))$, where $\mathrm{Diff}^+(\partial \mathrm{H_g})$ is the group of orientation preserving diffeomorphisms of ∂H_g. From here to the end of this paper, we assume $g \geq 2$.

The *cohomological dimension* of a group G, $cd(G)$, is defined to be the largest number n for which there exist a G-module M with $H^n(G, M)$ nonzero. We remark that if $G_1 \subset G_2$, then $cd(G_1) \leq cd(G_2)$. When G has torsion, $cd(G)$ is infinite. However, if G has finite index torsion free subgroups (we call G *virtually torsion free*), we define the *virtual cohomological dimension* of G, $vcd(G)$, to be the cohomological dimension of finite index torsion free subgroup $\hat{G}$. A theorem of Serre [11] states that this number is independent of the choice of $\hat{G}$. For vcd of $\mathcal{M}_g$ and $\mathcal{H}_g$, Harer [5] showed $vcd(\mathcal{M}_g) = 4g-5$, and McCullough [10] showed $vcd(\mathcal{H}_2) = 3$, and, if $g \geq 3$, $3g-2 \leq vcd(\mathcal{H}_g) \leq 4g-5$. In this paper, we will show the following result.

Theorem 1.1. *If $g \geq 2$, the virtual cohomological dimension of $\mathcal{H}_g$ is equal to $4g-5$.*

Date: February 16, 2001.

1991 *Mathematics Subject Classification.* 57N10, 57N05, 20F38.

Key words and phrases. virtual cohomological dimension, Euler number, 3-dimensional handlebody, mapping class group.

Hatcher observed that this result can be shown by investigation of the action of $\mathcal{H}_g$ on the disc complex defined by McCullough [10]. In this paper, we show this result with the explicit description of subgroup of $\mathcal{H}_g$ that achieves the $vcd(\mathcal{H}_g)$.

We give some remark on the relationship between $\mathcal{H}_g$ and the outer automorphism group of free group of rank g. We denote F_g the free group of rank g and $\mathrm{Out(F_g)}$ the outer automorphism group of it. There is a natural homomorphism from $\mathcal{H}_g$ to $\mathrm{Out(F_g)}$ defined by the action of diffeomorphisms on the fundamental group of H_g. This homomorphism is a surjection [4]. Culler and Vogtmann [3] showed that $vcd(\mathrm{Out(F_g)}) = 2g - 3$. This fact indicate that the kernel of the above surjection is, in some sense, big. In fact, McCullough [9] showed that the kernel of the above surjection is not finitely generated.

We review the Euler characteristics of groups (see [2]). For a group G of finite homological type and torsion-free, we define the *Euler characteristic* $\chi(G)$ by

$$\chi(G) = \sum_i (-1)^i \mathrm{rk}_{\mathbb{Z}}(H_i(G)).$$

For a group G of finite homological type which may have torsion, we choose a torsion free subgroup $\hat{G}$ of finite index, and define $\chi(G)$ by

$$\chi(G) = \frac{\chi(G)}{(G : \hat{G})},$$

where $(G : \hat{G})$ is the index of $\hat{G}$ in G. Since, $\mathcal{H}_g$ is of type VFL [10], we can define $\chi(\mathcal{H}_g)$. We will show the following result.

Theorem 1.2. $\chi(\mathcal{H}_g) = 0$.

2. Proof of Theorem 1.1.

In general, for an oriented C^∞-manifold A and its subset B, we denote $\mathrm{Diff^+(A)}$ the group of all orientation preserving diffeomorphisms of A, denote $\mathrm{Diff^+(A,\ fix\ B)}$ the group of elements of $\mathrm{Diff^+(A)}$ whose restriction to B is identity map, and denote $\mathrm{Diff^+(A,\ B)}$ the group of elements of $\mathrm{Diff^+(A)}$ which fixes B as a set. For a disc D in ∂H_g, we define $\mathcal{H}_g^1 = \pi_0(\mathrm{Diff^+(H_g, fix\ D)})$, and define $\mathcal{M}_g^1 = \pi_0(\mathrm{Diff^+(\partial H_g, fix\ D)})$. For the center p of the above disc D, we define $\mathcal{H}_{g,1} = \pi_0(\mathrm{Diff^+(H_g, fix\ \{p\})})$, and



FIGURE 1.

define $\mathcal{M}_{g,1} = \pi_0(\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix}\ \{p\}))$. Let $D_1, D_2, \ldots, D_g$ be the cocores of 1-handles which are used to construct H_g. These discs $D_1, D_2, \ldots, D_g$ are properly embedded discs in H_g. Let $E_1, \ldots, E_{g-1}$ and C be properly embedded discs as are indicated in Figure 1.

We introduce some elements of $\mathcal{H}_g$. For a disc D properly embedded in H_g, let N be a regular neighborhood of D in H_g. We parametrize N by $\phi : [-1, 1] \times D^2 \longrightarrow N$ such that, $\phi(\{0\} \times D^2) = D$, and $\phi([-1, 1] \times \partial D^2)$ is an annulus in ∂H_g. Let ψ be the diffeomorphisms of $[-1, 1] \times D^2$, defined by $\psi(t, r, \theta) = (t, r, \theta + (1 - t)\pi)$, where (r, θ) is a polar coordinate of D^2. The map δ_D from H_g to itself, defined by $\delta_D(x) = \phi \circ \psi \circ \phi^{-1}(x)$ iff $x \in N$, $= x$ iff $x \notin N$, is an orientation preserving diffeomorphism of H_g. We call this *disc twist* about D. The isotopy class of δ_D is an element of $\mathcal{H}_g$, and we call this *disc twist* about D and denote this d_D. For an annulus A properly embedded in H_g, let N be a regular neighborhood of A in H_g. We parametrize N by $\phi : [-1, 1] \times S^1 \times [0, 1] \longrightarrow N$, such that $\phi|_{\{0\} \times S^1 \times [0,1]}$ is a parametrization of A, $\phi([-1, 1] \times S^1 \times \{0\})$ and $\phi([-1, 1] \times S^1 \times \{1\})$ are disjoint annuli in ∂H_g. Let ψ be the diffeomorphism on $[-1, 1] \times S^1 \times [0, 1]$ defined by $\psi(t, \theta, s) = (t, \theta + (1 - t)\pi, s)$, where θ is a polar coordinate of S^1. The map α_A of H_g, defined by $\alpha_A(x) = \phi \circ \psi \circ \phi^{-1}(x)$ iff $x \in N$, $= x$ iff $x \notin N$, is an orientation-preserving diffeomorphism on H_g. We call this *annulus twist* about A. The isotopy class of α_A is an element of $\mathcal{H}_g$, and we call this *annulus twist* about A and denote this a_A.

We introduce some expressions used in this paper. Let N be a regular neighborhood of ∂H_g in H_g, and A be an annulus in ∂H_g. We parametrize N as $\phi : [0,1] \times \partial H_g \longrightarrow N$, such that $\phi(\{0\} \times \partial H_g) = \partial H_g$ and $\phi|_{\{0\} \times \partial H_g}$ is an identity map. The set $A' = \phi(\partial A \times [0,1] \cup A \times \{1\})$ is an annulus properly embedded in H_g. The sentence "we *push A into H_g*" means that we obtain A' from A. For a disc D in ∂H_g, the meaning of the sentence "we *push D into H_g*" is given in the same manner as above.

G. Mess [8] discovered some subgroups B_g, B_g^1 of the mapping class groups $\mathcal{M}_g$, $\mathcal{M}_g^1$ respectively. We call B_g and B_g^1 *Mess subgroups*. We review the definition of Mess subgroups (this definition is quoted from §6.3 of [7]). Mess subgroups are defined in a recursive manner.

Step 0: Let B_2 be the subgroup of $\mathcal{M}_2$ generated by Dehn twist about any three pairwise disjoint pairwise nonisotopic simple closed curves C_0, C_1, C_2 in ∂H_2.

Step 1_g: We assume that B_g $(g \geq 2)$ is already defined. There is a surjection from $\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix\ D})$ to $\mathrm{Diff}^+(\partial \mathrm{H_g})$ defined by forgetting the disc D, and this surjection induce an surjection $f : \mathcal{M}_g^1 \longrightarrow \mathcal{M}_g$. Let B_g^1 be the preimage of B_g under f.

Step 2_g: By the restriction of diffeomorphisms, we can define a homomorphism $\rho : \mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix\ D}) \longrightarrow \mathrm{Diff}^+(\partial \mathrm{H_g} - \mathrm{intD}, \mathrm{fix\ } \partial \mathrm{D})$. We consider some embedding $\partial H_g - \mathrm{intD}$ into ∂H_{g+1} and identify $\partial H_g - \mathrm{intD}$ with its image. The extension of diffeomorphisms of $\partial H_g - \mathrm{intD}$ fixed on ∂D by the identity across the complement of $\partial H_g - \mathrm{intD}$ in ∂H_{g+1} define a homomorphism $\iota : \mathrm{Diff}^+(\partial \mathrm{H_g} - \mathrm{intD}, \mathrm{fix} \partial \mathrm{D}) \longrightarrow \mathrm{Diff}^+(\partial \mathrm{H_{g+1}})$. The homomorphism $\iota \circ \rho$ induce a homomorphism $i : \mathcal{M}_g^1 \longrightarrow \mathcal{M}_{g+1}$. In the complement of $\partial H_g - \mathrm{intD}$ in ∂H_{g+1}, we choose some non-trivial simple close curve C and consider the Dehn twist $t \in \mathcal{M}_{g+1}$ about this curve. Let T be the infinite cyclic group generated by t. We define B_{g+1} as the group generated by $i(B_g^1)$ and T.

Mess showed the following theorem [8] (see also Corollary 6.3B of [7]).

Theorem 2.1. *The cohomological dimension of B_g is equal to $4g - 5$.* □

We will show the following lemma.

Lemma 2.2. *B_g is a subgroup of $\mathcal{H}_g$.*

Remark 2.3. The above fact is remarked by Mess [8, p.4].

The definition of B_g involves some choices. This lemma means that, with some good choices, B_g is realised as a subgroup of $\mathcal{H}_g$.

Proof. Along the steps of the definition of B_g, we will check that B_2, B_g^1, B_{g+1} can be constructed as subgroups of $\mathcal{H}_2$, $\mathcal{H}_g^1$, $\mathcal{H}_{g+1}$ respectively. In each steps, we use the same notations as used in definitions of B_g and B_g^1.

Step 0: We choose $C_0 = \partial D_1$, $C_1 = \partial C$, $C_2 = \partial D_2$, then $B_2 \subset \mathcal{H}_2$.

Step 1_g: We assume that $B_g \subset \mathcal{H}_g$. Let $g_1, \ldots, g_n$ be the generators of B_g. For each g_i, we can choose an element $\tilde{g}_i$ of $\mathcal{H}_g^1$ such that $f(\tilde{g}_i) = g_i$. By the definition, B_g^1 is generated by the kernel of f and $\tilde{g}_1, \ldots, \tilde{g}_n$. In order to obtain generators for the kernel of f, we consider the following two short exact sequences.

$$1 \longrightarrow \mathbb{Z} \longrightarrow \mathcal{M}_g^1 \xrightarrow{\alpha} \mathcal{M}_{g,1} \longrightarrow 1. \tag{S1}$$

$$1 \longrightarrow \pi_1(\partial H_g, \mathrm{fix}\{\mathrm{p}\}) \xrightarrow{\beta} \mathcal{M}_{\mathrm{g}}^1 \xrightarrow{\gamma} \mathcal{M}_{\mathrm{g}} \longrightarrow 1. \tag{S2}$$

The group $\mathbb{Z}$ in (S1) is an infinite cyclic group generated by the Dehn twist d about ∂D. The homomorphism α is induced from the homomorphism from $\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix\ D})$ to $\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix}\{\mathrm{p}\})$ defined by crushing D into a point p. The sequence (S2) is introduced by Birman [1]. The homomorphism γ is induced from the homomorphism from $\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix}\{\mathrm{p}\})$ to $\mathrm{Diff}^+(\partial \mathrm{H_g})$ defined by forgetting the point p. The group $\pi_0(\partial H_g, p)$ is generated by simple loops in ∂H_g, whose base points are p. Let $l_1, \ldots, l_{2g}$ be simple loops in ∂H_g, homotopy classes of which generates $\pi_0(\partial H_g, p)$. For each l_i, let L_i be an annulus in ∂H_g, which is a regular neighborhood of l_i such that $L_i \supset D \ni p$. ∂L_i is two simple closed curves l_i^1, l_i^2 in ∂H_g. The homomorphism β is defined so that it maps a homotopy class of l_i (denote $[l_i]$ for short) to a homotopy class of $\lambda_i = (+\text{Dehn twist about } l_i^1) \times (-\text{Dehn twist about } l_i^2)$. This homeomorphism λ_i is also an element of $\mathrm{Diff}^+(\partial \mathrm{H_g}, \mathrm{fix\ D})$, and $\alpha($ an element of $\mathcal{M}_g^1$ represented by $\lambda_i) = \beta([l_i])$. Let $\tilde{l}_i$ be an element of $\mathcal{M}_g^1$ represented by λ_i. Since the kernel of f is equal to α^{-1}(the kernel of γ) $= \alpha^{-1}$(the image of β), the kernel of f is generated by d and $\tilde{l}_1, \ldots, \tilde{l}_{2g}$. Let D' be a disc in H_g made by pushing D into H_g, and $\delta_{D'}$ be the disc twist about D'. Let L'_i be an annulus made by pushing L_i into H_g, and $\alpha_{L'_i}$ be the annulus twist about L'_i. The diffeomorphisms $\delta_{D'}$ and $\alpha_{L'_i}$ are elements of $\mathrm{Diff}^+(\mathrm{H_g}, \mathrm{fix\ D})$, and restrictions of them to ∂H_g represent d and $\tilde{l}_i$ respectively. This fact shows that the kernel of f is included in $\mathcal{H}_g^1$. Hence, $B_g^1 \subset \mathcal{H}_g^1$.

Step 2_g: It is easy to see that $i(B_g^1) \subset \mathcal{H}_{g+1}$. We choose $C = \partial D_{g+1}$, then $t \in \mathcal{H}_{g+1}$. Therefore, $B_{g+1} \subset \mathcal{H}_{g+1}$. ☐

Along the line of the proof of Theorem 6.4.A in [7], we will prove Theorem 1.1.

Proof of Theorem 1.1.

There is a natural homomorphism $\mathcal{M}_g \to Aut(\mathrm{H_1}(\partial \mathrm{H_g}, \mathbb{Z}/3\mathbb{Z}))$ defined by the action of diffeomorphisms on homology. Let Γ be the kernel of this homomorphism. By Ivanov [6, Corollary 1.5], Γ is torsion free. Therefore, Γ, $\mathcal{H}_g \cap \Gamma$, and $B_g \cap \Gamma$ are finite index torsion free subgroups of $\mathcal{M}_g$, $\mathcal{H}_g$, and B_g respectively. By the definition of virtual cohomological dimension, $vcd(\mathcal{M}_g) = cd(\Gamma)$, $vcd(\mathcal{H}_g) = cd(\mathcal{H}_g \cap \Gamma)$ and $vcd(B_g) = cd(B_g \cap \Gamma)$. By Harer [5, Theorem 4.1], $vcd(\mathcal{M}_g) = 4g - 5$, hence, $cd(\Gamma)$ $= 4g - g$. By Theorem 2.1, $vcd(B_g) = cd(B_g) = 4g - 5$, hence, $cd(B_g \cap \Gamma) = 4g - 5$. By Lemma 2.2, $B_g \cap \Gamma \subset \mathcal{H}_g \cap \Gamma \subset \Gamma$, therefore, $cd(B_g \cap \Gamma) \leq cd(\mathcal{H}_g \cap \Gamma) \leq cd(\Gamma)$. These fact show this theorem. ☐

3. PROOF OF THEOREM 1.2.

McCullough defined a *disc complex* L in [10] and used this to give an estimation for $vcd(\mathcal{H}_g)$. We review the definition of L. By a *disc* in H_g, we mean a properly embedded 2-disc in H_g. The disc is *essential* when ∂D does not bound a 2-disc in ∂H_g. The disc complex L of H_g is the simplicial complex whose vertices are the isotopy classes of essential discs in H_g, and whose simplices are defined by the rule that a collection of $n+1$ distinct vertices spans a n-simplex if and only if it admit a collection of representatives which are pairwise disjoint. McCullough showed the following Theorem.

Theorem 3.1. [10, Theorem 5.3] *The disc complex L of H_g is contractible.* $\square$

We use the following two Propositions about Euler characteristics of groups.

Proposition 3.2. [2, Proposition §IX 7.3(d)] *Let $1 \to G' \to G \to G'' \to 1$ be a short exact sequence of groups with G' and G'' of finite homology type. If G is virtually torsion free, then G is of finite homological type and $\chi(G) = \chi(G')\chi(G'')$.* $\square$

Proposition 3.3. [2, Proposition §IX 7.3(e')] *Let X be a contractible simplicial complex on which G act simplicially. For each simplex σ of X, let $G_\sigma = \{g \in G | g\sigma = \sigma\}$. If X has only finitely many cells mod G, and, for each simplex σ of X, G_σ is of finite homological type, then*

$$\chi(G) = \sum_{\sigma \in \mathcal{E}} (-1)^{\dim \sigma} \chi(G_\sigma),$$

where $\mathcal{E}$ is a set of representative for the cells of X mod G. $\square$

For each simplex $\sigma = < D_0, \dots, D_n >$ of L, $G_\sigma = \pi_0 \mathrm{Diff}^+(\mathrm{H_g},\ \mathrm{D_0} \cup \cdots \cup \mathrm{D_n})$. For the same simplex γ, let Γ_σ be the graph defined as follows. The vertices of Γ_σ correspond to the components of $H_g - D_0 \cup \cdots \cup D_n$. Each edge corresponds to one of $D_0, \dots, D_n$ and connects the vertices corresponding to the components attached along this disc. There is a natural homomorphism δ from G_σ to the group of automorphisms of Γ_σ. Let A_σ be the image of δ. Since the group of automorphisms of Γ_σ is a finite group, A_σ is a finite group. There are short exact sequences.

$$1 \longrightarrow \pi_0 \mathrm{Diff}^+(\mathrm{H_g},\ \mathrm{fix\ D_0} \cup \cdots \cup \mathrm{D_n}) \longrightarrow \mathrm{G}_\sigma \xrightarrow{\delta} \mathrm{A}_\sigma \longrightarrow 1. \tag{S3}$$

$$\begin{aligned} 1 \longrightarrow & \mathbb{Z}^{n+1} \longrightarrow \pi_0 \mathrm{Diff}^+(\mathrm{H_g},\ \mathrm{fix\ D_0} \cup \cdots \cup \mathrm{D_n}) \\ & \xrightarrow{\epsilon} \pi_0 \mathrm{Diff}^+(\mathrm{H_g/D_0} \cup \cdots \cup \mathrm{D_n},\ \mathrm{fix\ D_0/D_0} \cup \cdots \cup \mathrm{D_n/D_n}) \longrightarrow 1. \end{aligned} \tag{S4}$$

The homomorphism ϵ is induced by crashing each D_i into one point. The group $\mathbb{Z}^{n+1}$ is generated by disc twist about $D_0, \cdots, D_n$, and , as is well-known, $\chi(\mathbb{Z}^{n+1}) = \chi((S^1)^{n+1}) = 0$. By applying Proposition 3.2 to (S4), we see $\chi(\pi_0 \mathrm{Diff}^+(\mathrm{H_g},\ \mathrm{fix\ D_0} \cup \cdots \cup \mathrm{D_n})) = 0$. Therefore, by (S3), we obtain $\chi(G_\sigma) = 0$. Theorem 1.2 follows from the above observation, Theorem 3.1 and Proposition 3.3.

Acknowledgements

The author would like to express his gratitude to Prof. T. Akita and Prof. N. Kawazumi and Prof. D. McCullough for their helpful comments.

References

[1] J.S. Birman, Mapping class groups and their relationship to braid groups, *Com. Pure and App. Math.* 22(1969), 213–238

[2] K.S. Brown, *Cohomology of groups,* Graduate Texts in Math., Vol.87, Springer(1982)

[3] M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups, *Invent. Math.* 84(1986), 91–119

[4] H.B. Griffiths, Automorphisms of a 3-dimensional handlebody, *Abh. Math. Sem. Univ. Hamburg* 26(1964), 191–210

[5] J.L. Harer, The virtual cohomological dimension of the mapping class group of an orientable surface, *Invent. Math.* 84(1986), 157–176

[6] N.V. Ivanov, *Subgroups of Teichmüller modular groups,* Translations of Mathematical Monographs, Vol.115, AMS(1992)

[7] N.V. Ivanov, Mapping class groups, preprint(Dec.21, 1998)

[8] G. Mess, Unit tangent bundle subgroups of the mapping class groups, preprint(1990)

[9] D. McCullough, Twist groups of compact 3-manifolds, *Topology* 24(1985), 461–474

[10] D. McCullough, Virtually geometrically finite mapping class groups of 3-manifolds. *J. Diff. Geom.* 33(1991), 1–65

[11] J.-P. Serre, Cohomology des groupes discrets, In: Prospects in Mathematics. Ann. Math. Stud. 70 (1971), 77–169

Department of Mathematics, Faculty of Science and Engineering, Saga University, Saga, 840 Japan

E-mail address: `hirose@ms.saga-u.ac.jp`